

Recognizing & Treating Heart Failure

Scott Blois, M.D.
Boulder Heart
303-622-5979



1

Who is your speaker today?

- Born in North California. 53 years old
- Undergraduate college: UCLA (1984-1988)
- Medical School: Baylor College of Medicine, Houston, TX (1989-1993)
- Residency and Cardiology Training Univ. of Colorado Health Sciences Center, Denver (1993-2000)
- Cardiology practice Austin, TX. Head of Heart Failure clinic with 700 patients (2000-2015)
- Joined Boulder Heart August of 2015 (12 cardiologist group) to run HF program Sept 2015
- Boulder Heart HF clinic has greater than 700 patients seeing us with emphasis on co-management with patients, lots of HF teaching and education & state of art medicines and devices if needed.



2

Outline of Talk/Objectives

1. Understand definition of Heart Failure (HF).
2. Understand growing prevalence of HF and economic costs to USA.
3. Learn signs and symptoms of HF.
4. Learn some of leading causes of HF.
5. Learn tests to work up and diagnose suspected HF pts.
6. Briefly, hear about some of principle treatments of HF (medications, pacemakers, defibrillators).
7. Learn how CHF specialty clinic works and lifestyle risk factor modifications.
8. Questions.

3

Definition of Heart Failure (HF)


“... Specific term used to define the clinical syndrome which ensues when the heart is unable to pump enough blood to supply the metabolic needs of the body.”

- Bristow, Braunwald's Heart Disease textbook, 2000



4

Four Chambers of Heart: Importance of Left Ventricle in HF



5

Heart Failure-Epidemiology

- 1:5 people will develop HF during their lifetime.
- 5 million cases of symptomatic HF patients (estimated 5-6 million people with asymptomatic heart failure/heart weakening).
- 550,000 new cases of HF/ year in USA.
- 287,000 deaths from HF/year in USA.


6

Heart Failure: Economic Costs

- 1998: total HF costs \$20 billion
- 2014: total HF costs \$32 billion
- Estimated 2030: total HF costs \$72 billion
- Most costly heart disease in USA (more than heart attacks, bypass surgeries or heart stents).
- Average length of stay (LOS) for HF hospitalization 5 days. Average cost \$30-40 thousand/hospitalization.

7

Actual & Projected HF Population, 1950-2050



*Based on data from Phase I of the National Health and Nutrition Examination Survey III 1980 and U.S. Bureau of the Census data and projections appearing in Leonard Hayflick's "How and Why We Age," Balantine Books, New York, 1994.

Figure 1. Effect of the aging population on the prevalence of heart failure

8

What Are The Symptoms of Heart Failure?

Think **FACES**...

- **F**atigue
- **A**ctivities limited
- **C**hest congestion
- **E**dema or ankle swelling
- **S**hortness of breath



9

Signs/Symptoms of HF

Commonly mistaken diagnoses that eventually are found to be Heart Failure:

- Asthma
- Pneumonia
- Upper respiratory infection
- Depression and tiredness in the elderly

– Very important to diagnose HF early as this has a better treatment response and better prognosis.



10

Leading Causes of HF in USA

- **Coronary Artery Disease (CAD), post heart attack**
- **HTN (Uncontrolled High Blood Pressure)**
- **IDIOPATHIC (DOUBLE HIT THEORY)**
 - ? VIRAL EXPOSURE + GENETIC PREDISPOSITION
- **VALVULAR**
- **ALCOHOLIC**



11

Conditions That Preclude CHF

Table 1. Known Causes of Dilated Cardiomyopathy.

<p>Toxins</p> <p>Ethanol*</p> <p>Chemotherapeutic agents (doxorubicin, bleomycin)</p> <p>Cobalt*</p> <p>Antiretroviral agents (zidovudine,* didanosine,* zalcitabine*)</p> <p>Phenothiazines*</p> <p>Carbon monoxide*</p> <p>Lead*</p> <p>Cocaine*</p> <p>Mercury*</p> <p>Metabolic abnormalities</p> <p>Nutritional deficiencies (thiamine,* selenium,* carnitine*)</p> <p>Endocrinologic disorders (hypothyroidism,* acromegaly,* thyrotoxicosis,* Cushing's disease, pheochromocytoma,* diabetes mellitus)</p> <p>Electrolyte disturbances (hypocalcemia,* hypophosphatemia*)</p>	<p>Inflammatory or infectious causes</p> <p>Infectious</p> <p>Viral (coxsackie virus, cytomegalovirus,* human immunodeficiency virus)</p> <p>Rickettsial</p> <p>Bacterial (diphtheria*)</p> <p>Mycobacterial</p> <p>Fungal</p> <p>Parasitic (toxoplasmosis,* trichinosis, Chagas' disease)</p> <p>Noninfectious</p> <p>Collagen vascular disorders (scleroderma, lupus erythematosus, dermatomyositis)</p> <p>Hypersensitivity myocarditis*</p> <p>Sarcoidosis*</p> <p>Peripartum dysfunction*</p> <p>Neuromuscular causes</p> <p>Duchenne's muscular dystrophy</p> <p>Facioscapulohumeral muscular dystrophy</p> <p>Erb's limb-girdle dystrophy</p> <p>Myotonic dystrophy</p> <p>Friedreich's ataxia</p> <p>Familial cardiomyopathies</p>
--	---

12

Tests to order in suspected HF pt

- Electrocardiogram (ECG): often abnormal with hypertrophy of heart, evidence of previous heart attacks or slowed diseased electrical system.
- Labs to look for anemia, kidney, liver and thyroid function.
- Chest X ray to look for enlarged heart or water in lung fields.
- Echocardiogram (most important test)
- Brain Natriuretic Peptide Level (Blood test)



13

Diagnostic Testing: ECHO



14

Diagnosing HF

Lab test – “BNP” – brain natriuretic peptide

- Protein made in heart.
- Elevated if heart dilated and weakened and congested.
- Abnormal if >400pg/ml
- NT-pro BNP: slightly different assay. Abnormal if > 450.
- Good Sensitivity (90%), Very specific (98% negative predictive value)
- Other disease see BNP elevated: Pulmonary issues like COPD, acute pneumonia, acute or subacute blood clots in lungs (PE), chronic kidney insufficiency, age > 85 y/o, pulmonary hypertension.



15

Other heart tests maybe later to better define and make specific diagnosis:

- Heart catheterization “Coronary angiogram”: evaluate cholesterol blockages of heart arteries and take pressure readings in heart. Usually done thru artery and veins in wrist or groin.
- MRI of heart
- Stress test (Treadmill or Chemical stress)
- Labs
- Genetic Testing



16

New York Heart Association Functional Classification

- I No limitations
No symptoms with ordinary activities
- II Slight limitation
Symptoms with ordinary activities
- III Marked limitations
Symptoms with less than ordinary activities
- IV Symptoms of heart failure at rest



17

New Approach to the Classification of Heart Failure

Stage	Patient Description
A	High risk for developing heart failure (HF) <ul style="list-style-type: none"> ▪ Hypertension ▪ CAD ▪ Diabetes mellitus ▪ Family history of cardiomyopathy
B	Asymptomatic HF <ul style="list-style-type: none"> ▪ Previous MI ▪ LV systolic dysfunction ▪ Asymptomatic valvular disease
C	Symptomatic HF <ul style="list-style-type: none"> ▪ Known structural heart disease ▪ Shortness of breath and fatigue ▪ Reduced exercise tolerance
D	Refractory end-stage HF <ul style="list-style-type: none"> ▪ Marked symptoms at rest despite maximal medical therapy (eg, those who are recurrently hospitalized or cannot be safely discharged from the hospital without specialized interventions)

Carvedilol is indicated for use in patients with mild to severe chronic HF and in patients with HTN.
Hunt SA et al. J Am Coll Cardiol. 2001;38:2101-2113.

18

Table 7. Clinical Stages of Chronic Heart Failure (CHF) Associated with Systolic Dysfunction. Based on Symptoms and CHF Hospitalization Requirement

Stage	Description	NYHA Class	Annualized mortality %*	Hospitalizations/Year
A (Asymptomatic-mild)	Asymptomatic or only minimal symptoms, rare hospitalizations	I-II	2-5	<0.25
B (Mild-moderate)	Mild-moderate symptoms, infrequent hospitalizations	II-III	5-15	0.25-0.75
C (Advanced)	Moderate-severe symptoms, frequent hospitalizations	III-IV	15-25	.75-2
D (Severe)	Persistent severe symptoms, frequent prolonged or continuous hospitalizations	IV	>25	>2

19

Risk of HF and COVID-19

- Patients with chronic HF at risk of worse outcome if get COVID infection
- In general, 1% mortality if get COVID and no medical issues
- Patients with chronic HF who have gotten COVID infection, 10% mortality in registries.




20

That's the Bad News.... Now the Good News!!!

- Before 1990, if developed HF could treat with diuretics (Lasix), Digoxin and Morphine to get rid of congestion and swelling but couldn't get the heart stronger unless got heart transplant. Many studies <1990, five year survival with HF comparable to metastatic breast and lung cancers.
- In 1990's, research started to learn when one has Heart Failure, the body turns on 3-4 hormones (in heart, kidneys, adrenal gland, other organs) to make struggling heart work harder. Those hormones in long run are bad and toxic to heart. Treatment/ medications designed to block those hormone surges.



21



22

What are the Medications used for Heart Failure?

Experts recommend:

- Beta blockers - block high adrenalin levels in body. Can slow disease progression and make heart get stronger. (egs: Carvidilol "Coreg" or Metoprolol "Toprol")
- ACE inhibitors or ARBs: block renin-angiotension hormone surge from kidneys and adrenal glands. Can slow disease progression and make heart get stronger. Increases blood flow. (egs: Lisinopril, Monopril, Avapro, Diovan)
- Aldosterone blockers- block effects on high of elevated aldosterone hormone which causes heart to fibrosis, retain salt and water, weaken heart. Mild diuretic. (egs: Spironolactone "Aldactone", "Inspra")




23

Comparison of Crude, Annualized Mortality Rates with ACE-I's and Beta-blockers

Agents	NYHA Class	n	12-mo Placebo Mortality	12-mo Effect Size Decrease
Ace inhibitors	II-IV	7050	11%	16%
Beta-blockers	II-IV	8373	11%	36%
Combined Mortality Reduction		15423	11%	46%



24



25

ACEI and ARBs and COVID infection

- Initially some concern about taking ace inhibitor or Angiotension receptor blocker meds used in HF and treatment of hypertension might increase risk of getting COVID infection.
- SARS-CoV2 which causes COVID-19 binds to ACE2 receptor and gains entry into host cells. ACEI and ARBs upregulate ACE2 receptor expression. Hypothesized initially that taking these meds lead to more severe infections and adverse outcomes.
- Have found this is not true and stopping your ACEI or ARB if have chronic HF can be very dangerous.

Boulder Community Health

26

Medications Used in Heart Failure (II)

- Diuretics: helps urinate out water and salt. Controls symptoms of congestion (shortness of breath or swelling in legs). Examples: Furosemide "Lasix", Torsemide "Demadex", HCTZ
- Digoxin (used less frequently now unless have AFIB too): weakly helps heart beat stronger.
- Hydralazine/isordil: especially help in African-Americans with HF.

Boulder Community Health

27

Supplements in HF

- Overall not found to be very helpful.
- Omega three free fatty acids helpful. Helps prevent further heart attacks and dementia.
- Coenzyme Q-10..... If on statin.
- Multivitamin daily helpful.
- Iron if anemic.
- Rest of supplements not much if any benefit and overall trying to avoid polypharmacy and drug-drug reactions.

Boulder Community Health

28

New HF medications approved in 2015

- Medication: Entresto (Sacubitril/Valsartan)... a combination pill of Nephilysin inhibitor and angiotension inhibitor. Added another 20% reduction in mortality to HF patient on top of standard therapy. Currently being used in pts not tolerating ACEI/ARB or getting worse despite having been on ACEI/ARB. FDA approved July 2015.
- Medication: Corlaner (Ivabradine)... I channel inhibitor in electrical cells in heart slowly heart rate down. Has shown no mortality benefit but in patients with persistent high heart rates > 75 beats/minutes decreased chance of being hospitalized for HF. Only indicated in sinus rhythm pts not Afib.
- Both are new and brand only... Expensive but pharmacology companies are trying to give voucher to bring down cost.



29


Treatments – Pacemakers and Defibrillators

- Regular Pacemakers – if slow, diseased electrical system in heart.
- Biventricular Pacemakers – If long “QRS” on EKG
- Defibrillators – If LVEF <35% despite treatments at risk for serious, life-threatening heart arrhythmias. Recommendations are for defibrillators that can shock heart out of dangerous ventricular arrhythmia, if it occurs.



30

Advanced Therapies for HF: Mechanical Ventricular Assist Devices (VADS)



© MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED.

31

History of Transplant Medicine

- 1947: first human cadaveric kidney transplant done in USA (Boston).
- 1963: first liver transplant in USA (Denver).
- 1966: first pancreatic transplant done.
- 1967: first heart transplant done by Christian Barnard in South Africa on December 3. Patient survives 18 days.
- 1968: first heart transplant in USA by Norman Shumway at Stanford on January 6. Patient survives two weeks.



32

History of Transplant Medicine

- 1970: first heart-lung transplant.
- 1981: Cyclosporin discovered from mold in soil by German pharmacy company. Tested and approved for immunosuppression use in organ transplant recipients. Mortality and morbidity decreases dramatically. Heart transplant becomes more widespread thereafter.
- Now: 3,000-4,000 heart transplant recipients/year in USA.



33

Mortality

- 20,000 living cardiac transplant recipients currently in US.

Years Since Surgery	Survival
– 1y	85-90%
– 5yrs	65%
– 9yrs	50%
– 17yrs	25%



34

Risk Factor/ Disease Modification for HF Patients

- Patient Education on:
 - BP control (SBP 95-120. DBP <80)
 - fluid and salt restrictions (less than 2-4 grams Na/day)
 - pathophysiology of CHF and importance of hormones to help understand why polypharmacy needed. Helps with med compliance.
- Avoidance of NSAIDs
- Minimizing alcohol consumption
- Surveillance for cardiac ischemia
- Anticoagulation to reduce chance of TIA/ CVAs
- Sudden Death Prevention (holters, defibrillators)
- Prevention of Significant Anemia
- Immunizations
- Obesity/ Weight loss
- CHF Specialty Clinic referral options.


35

Risk factor modification: Flu/immunizations

- “Heart Failure in a Cold Climate: Seasonal Variations in HF-related Morbidity and Mortality,” JACC 2002.
 - Scotland 1990 to 1996
 - Elderly pts (> 75y/o), 15-18 % higher admission and mortality rate in winter months.
 - Author's conclusions:
 - Higher morbidity and mortality of HF pts attributable to concurrent respiratory illnesses.
 - Extra vigilance of HF pts advised in winter.
 - Immunize against pneumococcus and influenza.

36

Modification of Risk Factors: Seasonal Variation in Heart Failure Hospitalizations and Deaths



JACC, 2002

37

Risk factor modification: Obesity

- Update of Framingham study, NEJM 2002
 - Body Mass Index (BMI)
 - 18.5 to 24.9 normal
 - 25.0 to 29.9 overweight
 - > 30.0 obese
- Obesity linked to heart enlarging, HF and diabetes
- Men: 5% increase risk of developing HF per 1% increase in BMI
- Women: 7% increase risk of developing HF per 1% increase in BMI

38

Risk Factor Modification: NSAIDs

- “Association of NSAIDs with First Occurrence of HF and with Relapsing HF,” Arch Int Med 2002.
 - 7,277 pts followed in Rotterdam with HF.
 - People with HF who used NSAIDs on a regular daily basis had 10 fold risk of dying than if not using NSAIDs regularly.
 - Conclusions: 1) NSAIDs reduce renal perfusion; 2) impair adequacy of renal prostaglandin production. Renal prostaglandins crucial role in compensating renal hemodynamics in pts with LV systolic dysfunction; 3) increase serum creatinine levels.

39

Salt and Water Restrictions in Heart Failure

- Weakened heart has a difficult time pumping water and salt thru heart out to kidneys to urinate out of body.
- Many of deleterious hormones activated in HF cause water and salt/sodium retention.
- HF patient should limit total fluid intake per day to < 64 oz (6-8 glasses total fluid).
- HF patient should limit to sodium intake to < 1,800 mg per day



40

Smoking and Alcohol in Heart Failure

- Need to stop smoking completely. If continue smoking, poor prognosis even if on right HF medications and treatments. Heart is already weakened and continued cigarette/nicotine can cause constriction of blood vessels, new heart attacks and sticky blood that make HF patient at risk for stroke.
- Alcohol > two glasses per day turns into toxin to heart. Cut back heavy alcohol use or heart will not get better.



41

Exercise in Heart Failure

- Usually it is safe and important to exercise if have heart failure.
- Studies show MODERATE exercise 3-4x/week for 30 minutes each session has survival benefit in HF.
- Out patient cardiac rehab is now covered by Medicare for patients with chronic heart failure.



42

Understanding Need for Polypharmacy in Treating Heart Failure

- Many deleterious hormones turned on in HF (at least 3-4 hormone systems known already and more being discovered).
- No magic bullet pill that will cure HF.
- Usually need to take 4-7 meds to best treat HF. Luckily most of these are all now generic (\$4-10/month/medication).
- Need to learn to compulsively monitor daily weight with scale at home and diary.



43

Risk Factor Modification: CHF Referral Centers

Getting your care at a designated CHF clinic has many good outcomes:

- Get on best HF meds and at right doses.
- Many studies show better survival and less chance of getting hospitalized for HF decompensation if get care at designated HF center like Boulder Heart.
- More cost effective. Less expenses for both patient and healthcare system.
- More understanding of their disease process with more time for communication and explaining HF and answering questions.
- More accessibility to telephone RNs and answering questions.
- Outcomes at Boulder Foothills hospital for treating HF and survival one of best in state of Colorado and better than national norms.

44

Future Breakthroughs

- Genomic Testing – sample of blood to see that genes you have and what medications will work best for you. Are you a better responder to Toprol XL or Coreg?
- Stem Cell Replacement in Heart – Via surgical injection into injured part of heart or via blood transfusion
- Gene Replacement – to grow new muscle cells in heart
- Mechanical Hearts- continue to be smaller and less morbidity and last longer without clots in pump or pump parts wearing out.



45

Summary (I)

- Heart Failure (HF) is when the heart is not strong enough to pump blood to keep up with metabolic needs of body.
- HF is most common and expensive cardiac condition in USA. 1:5 people will develop HF in their life.
- Symptoms are many and vague. Shortness of breath, swelling body and fatigue most common.
- Echocardiogram (echo) is most important test to evaluate and confirm HF.



46

Summary (II)

- HF is not an irreversible disease usually. Since early 1990's, many medicines and pacemakers have been developed that can make HF a chronic, tolerable disease with potential for the heart muscle to regenerate and get stronger.
- Early detection of HF is important and portends to best chance of recovery and good overall prognosis.
- If one has HF, many lifestyle modifications are necessary to do well (moderate exercise, salt and water restrictions, adherence to complex polypharmacy regime, stop smoking, keep up with vaccinations, etc).
- HF clinics like the one we have at Boulder Heart are best at treating HF and usually have the best survival rates.



47

Questions?

Thank You!

Scott Blois, M.D.
Boulder Heart
303-622-5979



48